

The Backreacted Kähler Geometry of Wrapped Branes

Nakwoo Kim^{1,2,*}

¹*Department of Physics and Research Institute of Basic Science,
Kyung Hee University, Hoegi-dong,
Dongdaemun-gu, Seoul 130-701, Korea*

²*School of Natural Sciences, Institute for Advanced Study,
Princeton, NJ 08540, USA*

Abstract

For supersymmetric solutions of D3(M2) branes with $AdS_3(AdS_2)$ factor, it is known that the internal space is expressible as $U(1)$ fibration over Kähler space which satisfies a specific partial differential equation involving the Ricci tensor. In this paper we study the wrapped brane solutions of D3 and M2-branes which were originally constructed using gauged supergravity and uplifted to $D = 10$ and $D = 11$. We rewrite the solutions in canonical form, identify the backreacted Kähler geometry, and present a class of solutions which satisfy the Killing spinor equation.

PACS numbers: 11.25.-w, 04.65.+e

*Electronic address: nkim@khu.ac.kr, nkim@ias.edu

I. INTRODUCTION

According to the AdS/CFT correspondence [1], given a supersymmetric solution of $D = 10$ or $D = 11$ supergravity with an anti-de-Sitter (AdS) factor we expect there should be a dual supersymmetric conformal field theory (SCFT). The maximally supersymmetric solutions arise from M5, D3, or M2 branes in the near horizon limit. A rich class of nontrivial solutions with less supersymmetries can be obtained when such branes are put at the tip of a singular special-holonomy manifold. For 1/4-BPS and higher, the supergravity solutions are categorically written as a direct product of AdS and Sasaki-Einstein (SE) manifolds. The check of AdS/CFT duality involving a nontrivial SE manifold is a very actively studied research area these days. For IIB backgrounds $AdS_5 \times SE_5$, essential references are [2][3]. For $AdS_4 \times SE_7$ in M-theory, the identification of dual SCFT is not fully understood yet. The readers are referred to [4] for M2-branes in $\mathbb{C}^4/\mathbb{Z}_k$ orbifold, while for nontrivial SE backgrounds see e.g. [5] for a recent study.

A more sophisticated method to obtain SCFT and dual AdS is to consider branes wrapped on a supersymmetric cycle in a special holonomy manifolds. Via topological twisting, the lower-dimensional field theory on the brane can retain supersymmetry. The twisting is best facilitated in the framework of lower-dimensional gauged supergravity [6], and it is straightforward to uplift the explicit solutions to 10 or 11 dimensions. The dual field theories for wrapped branes have been mysterious for many years, especially for M5 and M2-branes. For M5-branes the $D = 4$ SCFTs constructed in [7] may provide the answer. It is illustrated there that M5-branes wrapped on a Riemann surface with punctures give rise to a variety of novel quiver gauge theories which flow to $\mathcal{N} = 2$ SCFT in IR. It is argued in [8] that the gravity dual in the large- N limit should be $\mathcal{N} = 2$ solutions of [6], by comparing the central charges or the moduli space of marginal operators.

On the other hand, one may systematically explore the consequence of unbroken supersymmetry and AdS geometry in supergravity solutions. Typically one starts with the existence of Killing spinor fields, and look for the form of supergravity solutions by analyzing the algebraic and differential identities between spinor bilinears. This method has been especially powerful with 1/2-BPS geometries [9][10]. For 1/4-BPS and below the form of the metric is rather involved, but with the use of ingenious ansatz one may sometimes find new solutions. In particular, inhomogeneous SE manifolds $Y^{p,q}$ were found this way [11][12].

Some years ago the author applied this Killing spinor technique to the case of D3-branes geometry with an AdS_3 factor [13]. More concretely, we look for AdS_3 solutions in IIB supergravity where only the metric and five-form field strength are turned on. The result is quite elegant: the seven-dimensional internal space takes the form of twisted $U(1)$ fibration over six-dimensional Kähler space satisfying a higher-order differential equation involving Ricci tensor and scalar curvature.

$$\square R - \frac{1}{2}R^2 + R_{ij}R^{ij} = 0. \quad (1)$$

The warp factor for AdS_3 and the twisting of $U(1)$ can be fixed once the Kähler space is determined.

Remarkably, one discovers basically the same structure for M2-brane configurations containing AdS_2 [14]. This time the internal space is $U(1)$ fibration over eight-dimensional space satisfying again (1). It was illustrated that many known solutions with AdS_3 or AdS_2 factor can be rewritten this way, and also some new AdS_3, AdS_2 solutions were constructed by solving (1) directly [15–18]. (See also [19–22] for general properties of wrapped D3 and M2-branes.) It is also shown that this structure can be generalized to arbitrary higher dimensions [17]: $U(1)$ fibration over a solution of (1) can be thought of a solution of a $(2n + 1)$ -dimensional bosonic action comprised of metric, one-form and a scalar field. Existence of Killing spinor equation is reduced to (1) for any $n \geq 3$. It will be also worthwhile to comment that (1) was recently reproduced as a special case of extremal black hole geometry in [23].

In this paper we are interested in AdS_3 and AdS_2 solutions of Refs. [6] and [24] which are respectively constructed from $D = 5$ and $D = 4$ gauged supergravity theories and interpreted as D3 and M2 branes wrapped on Riemann surface. We will first identify the six and eight-dimensional Kähler subspace which satisfy (1), from the wrapped brane solutions in [6, 24]. For the case of D3 wrapped on H^2 in Calabi-Yau 3-manifold (CY3), it turns out that the six-dimensional space is locally $H^2 \times S^2$ fibred over a squashed S^2 . Similarly, M2-branes wrapped on a H^2 2-cycle in CY4 lead to a solution of (1) which is locally $H^2 \times KE_{2n}$ over a squashed S^2 (KE_{2n}^+ is Kähler-Einstein manifold with real dimension $2n$ and positive curvature). Motivated by these solutions, we consider a general ansatz for (1) and find solutions which are locally $H^2 \times KE_{2n}^+$ over squashed S^2 , which is our main result in this article.

This paper is organized as follows. In section II we review the Killing spinor geometry of [17] and explain how the solutions can be embedded in IIB and M-theory. In section III, we consider the wrapped D3-brane solution of [6] and M2-brane solution in [24] and identify the Kähler space satisfying (1). In section IV we consider a higher dimensional generalization which takes a form of $H^2 \times KE_{2n}^+$ over S^2 . We conclude in section V.

II. BPS GEOMETRY OF AdS_3 FROM D3, AND AdS_2 FROM M2

Here we review the geometries studied in [13–15] and generalized in [17]. We are interested in the following Lagrangian density in $(2n + 1)$ -dimensions ($n \geq 3$),

$$\mathcal{L} = e^{(1-n)B} \left[R + \frac{n(2n-3)}{2} (\nabla B)^2 + \frac{1}{4} e^{2B} F_{ab} F^{ab} - \frac{2n}{(n-2)^2} \right]. \quad (2)$$

The field contents consist of the metric, a scalar B , and a vector field strength $F = dA$. We may consider the same action defined for a space with Lorentzian signature, but for our purpose here it is understood that the metric defines a Riemannian manifold.

It is shown in [17] that the above Lagrangian admits a particular type of Killing spinors. More concretely, we consider Killing spinor equations

$$\begin{aligned} \left[\Gamma^a \nabla_a B + i \frac{2(n-1)}{n-2} + \frac{1}{2} e^B F_{ab} \Gamma^{ab} \right] \eta &= 0, \\ \left[\nabla_c + \frac{i}{2} \Gamma_c + \frac{1}{8} e^B F_{ab} \Gamma_c^{ab} \right] \eta &= 0. \end{aligned} \quad (3)$$

η is a spinor field in $(2n+1)$ -dimensions and Γ_a 's are generators of $\text{Cliff}(2n+1)$. As explained in [15], one can check that the existence of Killing spinor η , the Bianchi identity $dF = 0$, and the gauge field equation $d(e^{(3-n)B} * F) = 0$ guarantee that the equations of motion derived from (2) are satisfied.

Furthermore, the above models describe in fact interesting subsectors of $D = 10$ IIB supergravity and $D = 11$ supergravity, for $n = 3$ and $n = 4$ respectively. Given a Killing spinor solution to (3) for $n = 3$, we can embed it in a $D = 10$ IIB solution as follows:

$$\begin{aligned} ds^2 &= e^{-B/2} [ds^2(AdS_3) + ds_7^2], \\ F_5 &= -\frac{1}{4} [\text{Vol}(AdS_3) \wedge F - *_7 F]. \end{aligned} \quad (4)$$

We have a similar mapping for $n = 4$ case, with M-theory.

$$\begin{aligned} ds^2 &= e^{-2B/3} [ds^2(AdS_2) + ds_9^2] , \\ G_4 &= \text{Vol}(AdS_2) \wedge F. \end{aligned} \tag{5}$$

And of course the Killing spinor equations can be directly derived from the field ansatz and supersymmetry transformation rules of $D = 10$ and $D = 11$ supergravity.

One can proceed further and study the consequence of Killing spinors on the form of metric and other fields. It is a standard Killing spinor technique which has been extensively studied in recent literature. It turns out that our $(2n+1)$ -dimensional solution can be always written locally

$$ds_{2n+1}^2 = c^2(dz + P)^2 + e^B ds_{2n}^2 \tag{6}$$

i.e. as $U(1)$ vibration over $(2n)$ -dimensional space. Here $c = n/2 - 1$, ∂_z is a Killing vector and the $(2n+1)$ -dimensional fields are independent of z . Most importantly it turns out that the $(2n)$ -dimensional base space should be Kähler, and satisfy a *master equation*

$$\square R - \frac{1}{2}R^2 + R_{ij}R^{ij} = 0. \tag{7}$$

Here P is a one-form and related to Kähler geometry such that dP is the Ricci two-form for ds_{2n}^2 . The scalar field and the gauge field are then completely determined by the geometric data, as

$$\begin{aligned} e^B &= c^2 \left(\frac{R}{2} \right), \\ F &= -\frac{1}{c}J + c d [e^{-B}(dz + P)] . \end{aligned} \tag{8}$$

where J is the Kähler two-form of ds_{2n}^2 .

As discussed in [17] the physical application is not clear for $n > 4$, but it is mathematically intriguing that there exists a class of *pseudo-supergravity* models for all $n \geq 3$ which includes physically interesting solutions for String/M-theory.

III. WRAPPED BRANE SOLUTIONS FOR D3 IN CY3, AND M2 IN CY4

A. D3-branes wrapping a 2-cycle in CY3

We will start with a specific $D = 10$ IIB solution presented as D3-branes wrapped on a 2-cycle in Calabi-Yau 3-manifold, in [6]. The solution is best constructed in $D = 7$ gauged

supergravity first, and then uplifted to $D = 10$. In $D = 7$, the metric ansatz contains H^2 part and we also turn on magnetic flux on it. When acting on spinor fields, the spin connection of H^2 effectively cancel against the nonzero magnetic field, and this mechanism through AdS/CFT implies that that we have topological twist of boundary field theory.

As given in Eq.(18) of [6], the ten-dimensional metric is

$$2^{3/2}ds^2 = \sqrt{\Delta}ds_{AdS_3}^2 + \frac{1}{\sqrt{\Delta}} \left[\Delta ds_{H^2}^2 + 2\Delta d\theta^2 + 4\cos^2\theta(d\psi^2 + \sin^2\psi D\phi_1^2 + \cos^2\psi D\phi_2^2) + 2\sin^2\theta d\phi_3^2 \right], \quad (9)$$

where $\Delta = 1 + \sin^2\theta$. In this subsection we denote $L^2 = 2^{-3/2}$. In our convention H^2 has unit radius, and $d(D\phi_1) = d(D\phi_2) = -\frac{1}{2}\text{Vol}(H^2)$.

When we compare this metric with (4) and (6), it is obvious that $e^{-B} = L^4\Delta$ and the seven-dimensional internal space is to be rewritten as

$$\begin{aligned} & \frac{1}{\Delta} \left[\Delta ds_{H^2}^2 + 2\Delta d\theta^2 + 4\cos^2\theta(d\psi^2 + \sin^2\psi D\phi_1^2 + \cos^2\psi D\phi_2^2) + 2\sin^2\theta d\phi_3^2 \right] \\ &= \frac{1}{4}(dz + P)^2 + \frac{1}{L^4\Delta}ds_6^2. \end{aligned} \quad (10)$$

The constant-norm Killing vector ∂_z must be a linear combination of ∂_{ϕ_i} , ($i = 1, 2, 3$). We find it convenient to introduce a new set of coordinates as $\phi = \phi_1 - \phi_2$, $\varphi = \phi_1 + \phi_2$, $\chi = 2\psi$ and $z = 2\phi_3$. The internal space metric and the twisting one-form P are then found as

$$\begin{aligned} \frac{1}{L^4}ds_6^2 &= \Delta ds^2(H^2) + 2\Delta d\theta^2 + \frac{\sin^2 2\theta}{2\Delta}(d\varphi - \cos\chi d\phi)^2 + \cos^2\theta(d\chi^2 + \sin^2\chi d\phi^2) \quad (11) \\ P &= \frac{2}{\Delta}(d\varphi - \cos\chi\phi). \quad (12) \end{aligned}$$

Now what we should check is that the $d = 6$ metric in (11) is Kähler and satisfies (7) and dP is the Ricci two-form. This will be shown as a special case of a general construction in section IV.

B. AdS_2 from wrapped M2 branes

The prescription for supergravity solutions of wrapped branes [6] was generalized to M2-branes in [24]. It is convenient to consider a consistent subsector of $D = 4, \mathcal{N} = 8$ supergravity with $U(1)^4$, and study $AdS_2 \times H^2$ ansatz with magnetic flux on H^2 . The

solution we are interested in corresponds to M2-branes wrapped on a 2-cycle in CY 4-manifold presented in Eq.(3.19) of [24]. Using the uplifting formula (see for instance Eq.(2.1) in [24]), it is easy to check that the $D = 11$ metric is

$$ds_{11}^2 = \frac{\tilde{\Delta}^{2/3}}{2 \cdot 3^{5/3} \cdot e^2} \left[ds^2(AdS_2) + 3ds^2(H_2) + \frac{12}{\tilde{\Delta}} \left(3 \sum_{\alpha=1}^3 (d\mu_\alpha^2 + \mu_\alpha^2 D\phi_\alpha^2) + (d\mu_4^2 + \mu_4^2 d\phi_4^2) \right) \right]. \quad (13)$$

Here μ_a satisfy $\sum_{a=1}^4 \mu_a^2 = 1$ and together with unconstrained angles ϕ_a ($a = 1, \dots, 4$) they parametrize S^7 . The warp factor is $\tilde{\Delta} = \mu_1^2 + \mu_2^2 + \mu_3^2 + 3\mu_4^2$ and the coupling constant e^2 is related to the $D = 4$ cosmological constant as $\Lambda_4 = -12e^2$. The twisted one-forms satisfy $d(D\phi_\alpha) = -\frac{1}{3}\text{Vol}(H^2)$ for $\alpha = 1, 2, 3$.

To rewrite this metric in the form (5), we first note that $e^{-B} = L^3 \tilde{\Delta}$ with $L^{-2} = 2 \cdot 3^{5/3} \cdot e^2$. And in order to identify the constant norm Killing vector, we re-define some of angular coordinates as

$$\phi_a \rightarrow \phi_a + \phi_4/3, \quad a = 1, 2, 3. \quad (14)$$

Then one may easily check that the metric is written as (5) with

$$P = \frac{6}{\tilde{\Delta}} \sum_{a=1}^3 \mu_a^2 D\phi_a \quad (15)$$

and the remaining $d = 8$ Kähler base space is

$$\begin{aligned} \frac{1}{L^3} ds_8^2 &= 3\tilde{\Delta} ds^2(H_2) + 12 \left(3 \sum_{a=1}^3 d\mu_a^2 + d\mu_4^2 \right) \\ &\quad + 36 \sum_{a=1}^3 \mu_a^2 D\phi_a^2 - \frac{36}{\tilde{\Delta}} \left(\sum_{a=1}^3 \mu_a^2 D\phi_a \right)^2 \end{aligned} \quad (16)$$

It still looks complicated, but in fact one can rewrite this metric as $H^2 \times \mathbb{CP}^2$ over squashed S^2 , i.e. a natural generalization of (11). Let us define

$$\mu_4 = \sin \theta, \quad \mu_a = \tilde{\mu}_a \cos \theta \quad (a = 1, 2, 3). \quad (17)$$

And for ϕ_a angles we single out the diagonal part and define

$$\phi_a = \tilde{\phi}_a + \psi \quad (a = 1, 2, 3). \quad (18)$$

Note that $\sum_{a=1}^3 \tilde{\mu}_a^2 = 1$ and $\sum_{a=1}^3 \tilde{\phi}_a = 0$. The Ricci potential is now

$$P = \frac{6 \cos^2 \theta}{\tilde{\Delta}} \left(D\psi + \sum_{a=1}^3 \tilde{\mu}_a^2 d\tilde{\phi}_a \right), \quad d(D\psi) = -\text{Vol}(H^2). \quad (19)$$

The metric becomes

$$\begin{aligned} \frac{1}{L^3} ds_8^2 = & 3\tilde{\Delta} ds^2(H_2) + 12\tilde{\Delta} d\theta^2 + \frac{27 \sin^2 2\theta}{\tilde{\Delta}} (d\psi + \sum_{a=1}^3 \tilde{\mu}_a^2 d\tilde{\phi}_a)^2 \\ & + 36 \cos^2 \theta \left(\sum_{a=1}^3 d\tilde{\mu}_a^2 + \sum_{a=1}^3 \tilde{\mu}_a^2 d\tilde{\phi}_a^2 - \left(\sum_{a=1}^3 \tilde{\mu}_a^2 d\tilde{\phi}_a \right)^2 \right). \end{aligned} \quad (20)$$

We see that the part of the metric within big parentheses in the second line gives $\mathbb{CP}^2$. The easiest way to check this fact is to start with flat $\mathbb{C}^n$, parametrized by $z_j = \tilde{\mu}_j e^{i\tilde{\phi}_j}$ and rewrite the metric as metric cone of S^{2n-1} , which in turn is written as $U(1)$ fibration over $\mathbb{CP}^{n-1}$.

The remaining task of checking (7) etc will be relegated to section IV, just as wrapped D3-brane case.

IV. GENERAL ANSATZ AND SOLUTIONS

Motivated by the explicit solutions in section III, we consider the following metric ansatz for (7).

$$ds_{2n+4}^2 = \Delta_1 ds^2(H_2) + \Delta_2 d\theta^2 + \frac{\sin^2 2\theta}{\Delta_2} D\psi^2 + \cos^2 \theta ds^2(KE_{2n}^+), \quad (21)$$

where Δ_1, Δ_2 are functions of θ . For $n = 1, 2$ we expect to reproduce the wrapped D3 and M2-brane solutions explained in section III. The nontrivial $U(1)$ -fibration is defined as

$$D\psi = d\psi - P_2 - P_{2n}. \quad (22)$$

Here P_2 is Ricci potential for H_2 , and P_{2n} is the Ricci potential for KE_{2n}^+ . We assume the overall normalization is such that

$$dP_2 = -J_2, \quad dP_{2n} = +J_{2n}. \quad (23)$$

In other words,

$$d(D\psi) = J_2 - J_{2n}. \quad (24)$$

To check the Kähler condition, we define the complex structure so that the Kähler 2-form is

$$J_{2n+4} = \Delta_1 J_2 + \sin 2\theta d\theta \wedge D\psi + \cos^2 \theta J_{2n}. \quad (25)$$

One can easily check J_{2n+4} is closed if $\Delta'_1 = \sin 2\theta$, i.e.

$$\Delta_1 = c + \sin^2 \theta \quad (26)$$

where c is constant.

Now let us consider $(n, 0)$ -form Ω . As it is well-known, we calculate $d\Omega$ and if it can be written as $d\Omega = iP \wedge \Omega$, the metric is Kähler with dP the Ricci 2-form. We have

$$\Omega = e^{-i\psi} \Omega_2 \wedge \cos^n \theta \left(\sqrt{\Delta_1 \Delta_2} d\theta + i \sqrt{\frac{\Delta_1}{\Delta_2}} \sin 2\theta D\psi \right) \wedge \Omega_{2n} \quad (27)$$

where we inserted the phase factor $e^{i\psi}$ for later convenience. This coefficient of course does not affect the result for Ricci-tensor. It is straightforward to check

$$d\Omega_{2n+4} = iP_{2n+4} \wedge \Omega_{2n+4}, \quad (28)$$

$$P_{2n+4} = - \left[1 + \frac{1}{\cos^n \theta \sqrt{\Delta_1 \Delta_2}} \frac{d}{d\theta} \left(\sqrt{\frac{\Delta_1}{\Delta_2}} \sin 2\theta \cos^n \theta \right) \right] D\psi \equiv f D\psi. \quad (29)$$

Summarizing, we have established that the ansatz (21) always gives us a locally Kähler space for any $\Delta_1 = c + \sin^2 \theta$ and arbitrary $\Delta_2(\theta)$.

We can then compute the Ricci tensor by considering $\mathcal{R}_{2n+4} = dP_{2n+4}$. The computation is rather involved for general Δ_2 , since (7) is nonlinear and involves a fourth derivative. We have not been able to find the most general solution, but if we assume Δ_2 is also a polynomial of $\sin^2 \theta$, it is not difficult to check that the following is the only possibility for regular solutions:

$$\Delta_2 = 2n\Delta_1 = 2(n \sin^2 \theta + 1). \quad (30)$$

For reference, we record that

$$f = -\frac{2 \cos^2 \theta}{1 + n \sin^2 \theta}, \quad R = \frac{4(1 + n)}{1 + n \sin^2 \theta}. \quad (31)$$

For $n = 1$ this solution reduces to wrapped D3-branes, and for $n = 2$ and $KE_4 = \mathbb{CP}^2$ corresponds to M2 solutions presented in section III. We can also take $KE_4 = S^2 \times S^2$ and this gives us a physically distinct solution for a M2-brane configuration. It will be also interesting to study the field theory dual of this new AdS_2 solution.

V. DISCUSSION

In this paper we have chosen the wrapped brane solutions constructed using gauged supergravity in [6][24], and identified the back reacted Kähler geometry in terms of the general description in [17]. The main motivation of this work was to obtain new AdS solutions using the generalized ansatz (21) which were drawn from wrapped brane solutions. By directly solving the master equation (7), we were able to find the natural higher-dimensional generalization.

The advantage of (7) is its generality, so in principle one can find any solution which probably cannot be obtained from lower-dimensional gauged supergravity system. In particular one can slightly change the ansatz of (21) and look for solutions. Consider the following metric ansatz

$$ds_{2n+4}^2 = \Delta_1 ds^2(K E_{2n}^-) + \Delta_2 d\theta^2 + \frac{\sin^2 2\theta}{\Delta_2} D\psi^2 + \cos^2 \theta (d\chi^2 + \sin^2 \chi d\phi^2) \quad (32)$$

where

$$D\psi = d\psi + \cos \chi d\phi - P_{2n}, \quad dP_{2n} = -J_{2n}. \quad (33)$$

For instance the space can be chosen as $\otimes_{i=1}^n H_i^2 \times S^2$ over squashed S^2 . Now that we have increased the dimensionality of negatively curved part of the metric and we do not expect this class of metric may be covered by gauged supergravity. We have checked that this space allows Kähler structure provided $\Delta_1 = a + \sin^2 \theta$, and computed Ricci tensor etc. Eq.(7) is again quite a complicated expression of Δ_2 but one can proceed by assuming Δ_2 is also a polynomial of $\sin^2 \theta$. Unfortunately we have not found any physically interesting solution for $n > 1$. We noticed for instance that $\Delta_1 = \sin^2 \theta - \frac{3}{7}$, $\Delta_2 = 6\Delta_1$ locally satisfies (7) for $n = 3$, but this space is singular since the hyperbolic space collapses at $\sin^2 \theta = 3/7$.

Recently there is a revival of interest in the gravity solution of wrapped brane solutions. The authors of [25] considered a generalization of the gauged supergravity ansatz of [6] where the Riemann surface 2-cycle starts with an arbitrary metric in UV. The motivation of [25] was to prove a crucial assumption of [7] that for wrapped M5-branes the conformal factor of 2-cycle should be irrelevant in IR. This can be seen indeed from the elliptic flow equation derived in [25]. But we would like to point out the central result of [25] was derived using $D = 7$ gauged supergravity, not the original $D = 11$ supergravity. It will be certainly desirable to study more general form of supersymmetric solutions for wrapped M5-branes

(see e.g. [26]) and study the flow equation, directly in $D = 11$. On the other hand in [27, 28] new AdS_5 solutions and their field theory duals are constructed using general $\mathbb{C}^2$ bundle over Riemann surface. If this construction can be also applied to D3 or M2 branes it will be interesting to identify the corresponding Kähler solution to (7).

Acknowledgments

This work is supported by the sabbatical leave petition program (2012) of Kyung Hee University, the National Research Foundation of Korea (NRF) funded by the Korean Government (MEST) with grant No. 2009-0085995, 2010-0023121, and also through the Center for Quantum Spacetime (CQUeST) of Sogang University with grant No. 2005-0049409.

-
- [1] J. M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, *Adv.Theor.Math.Phys.* **2** (1998) 231–252, [[hep-th/9711200](#)].
 - [2] I. R. Klebanov and E. Witten, *Superconformal field theory on three-branes at a Calabi-Yau singularity*, *Nucl.Phys.* **B536** (1998) 199–218, [[hep-th/9807080](#)].
 - [3] D. Martelli and J. Sparks, *Toric geometry, Sasaki-Einstein manifolds and a new infinite class of AdS/CFT duals*, *Commun.Math.Phys.* **262** (2006) 51–89, [[hep-th/0411238](#)].
 - [4] O. Aharony, O. Bergman, D. L. Jafferis, and J. Maldacena, *$N=6$ superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*, *JHEP* **0810** (2008) 091, [[arXiv:0806.1218](#)].
 - [5] C. Closset and S. Cremonesi, *Toric Fano varieties and Chern-Simons quivers*, [arXiv:1201.2431](#).
 - [6] J. M. Maldacena and C. Nunez, *Supergravity description of field theories on curved manifolds and a no go theorem*, *Int. J. Mod. Phys.* **A16** (2001) 822–855, [[hep-th/0007018](#)].
 - [7] D. Gaiotto, *$N=2$ dualities*, [arXiv:0904.2715](#).
 - [8] D. Gaiotto and J. Maldacena, *The gravity duals of $N=2$ superconformal field theories*, [arXiv:0904.4466](#).
 - [9] H. Lin, O. Lunin, and J. M. Maldacena, *Bubbling AdS space and $1/2$ BPS geometries*, *JHEP* **10** (2004) 025, [[hep-th/0409174](#)].

- [10] E. O Colgain, J.-B. Wu, and H. Yavartanoo, *On the generality of the LLM geometries in M-theory*, *JHEP* **1104** (2011) 002, [[arXiv:1010.5982](#)].
- [11] J. P. Gauntlett, D. Martelli, J. Sparks, and D. Waldram, *Supersymmetric AdS(5) solutions of M theory*, *Class.Quant.Grav.* **21** (2004) 4335–4366, [[hep-th/0402153](#)].
- [12] J. P. Gauntlett, D. Martelli, J. Sparks, and D. Waldram, *Sasaki-Einstein metrics on $S^{*2} \times S^{*3}$* , *Adv.Theor.Math.Phys.* **8** (2004) 711–734, [[hep-th/0403002](#)].
- [13] N. Kim, *AdS(3) solutions of IIB supergravity from D3-branes*, *JHEP* **0601** (2006) 094, [[hep-th/0511029](#)].
- [14] N. Kim and J.-D. Park, *Comments on AdS(2) solutions of D=11 supergravity*, *JHEP* **0609** (2006) 041, [[hep-th/0607093](#)].
- [15] J. P. Gauntlett, N. Kim, and D. Waldram, *Supersymmetric AdS(3), AdS(2) and Bubble Solutions*, *JHEP* **0704** (2007) 005, [[hep-th/0612253](#)].
- [16] H. Kim, K. K. Kim, and N. Kim, *1/4-BPS M-theory bubbles with $SO(3) \times SO(4)$ symmetry*, *JHEP* **0708** (2007) 050, [[arXiv:0706.2042](#)].
- [17] J. P. Gauntlett and N. Kim, *Geometries with Killing Spinors and Supersymmetric AdS Solutions*, *Commun.Math.Phys.* **284** (2008) 897–918, [[arXiv:0710.2590](#)].
- [18] A. Donos, J. P. Gauntlett, and N. Kim, *AdS Solutions Through Transgression*, *JHEP* **0809** (2008) 021, [[arXiv:0807.4375](#)].
- [19] O. A. Mac Conamhna and E. O Colgain, *Supersymmetric wrapped membranes, AdS(2) spaces, and bubbling geometries*, *JHEP* **0703** (2007) 115, [[hep-th/0612196](#)].
- [20] P. Figueras, O. A. Mac Conamhna, and E. O Colgain, *Global geometry of the supersymmetric AdS(3)/CFT(2) correspondence in M-theory*, *Phys.Rev.* **D76** (2007) 046007, [[hep-th/0703275](#)].
- [21] J. P. Gauntlett and O. A. Mac Conamhna, *AdS spacetimes from wrapped D3-branes*, *Class.Quant.Grav.* **24** (2007) 6267–6286, [[arXiv:0707.3105](#)].
- [22] E. O Colgain, J.-B. Wu, and H. Yavartanoo, *Supersymmetric AdS3 X S2 M-theory geometries with fluxes*, *JHEP* **1008** (2010) 114, [[arXiv:1005.4527](#)].
- [23] U. Gran, J. Gutowski, and G. Papadopoulos, *AdS backgrounds from black hole horizons*, [arXiv:1110.0479](#).
- [24] J. P. Gauntlett, N. Kim, S. Pakis, and D. Waldram, *Membranes wrapped on holomorphic curves*, *Phys. Rev.* **D65** (2002) 026003, [[hep-th/0105250](#)].

- [25] M. T. Anderson, C. Beem, N. Bobev, and L. Rastelli, *Holographic Uniformization*,
`arXiv:1109.3724`.
- [26] J. P. Gauntlett, O. A. Mac Conamhna, T. Mateos, and D. Waldram, *AdS spacetimes from wrapped M5 branes*, *JHEP* **0611** (2006) 053, [`hep-th/0605146`].
- [27] I. Bah, C. Beem, N. Bobev, and B. Wecht, *AdS/CFT Dual Pairs from M5-Branes on Riemann Surfaces*, `arXiv:1112.5487`.
- [28] I. Bah, C. Beem, N. Bobev, and B. Wecht, *Four-Dimensional SCFTs from M5-Branes*, *JHEP* **06** (2012) 005, [`arXiv:1203.0303`].